

SURAT KETERANGAN

Nomor: 1093/UNUSA/Adm-LPPM/X/2020

Lembaga Penelitian dan Pengabdian Kepada Masyarakat (LPPM) Universitas Nahdlatul Ulama Surabaya menerangkan telah selesai melakukan pemeriksaan duplikasi dengan membandingkan artikel-artikel lain menggunakan perangkat lunak **Turnitin** pada tanggal 20 Oktober 2020.

Judul : Monthly Water Balance Analysis for the Determination of Available Growing Season at North Samarinda District, East Kalimantan
Penulis : A.Y Pulihasih, A.P Sujalu
No. Pemeriksaan : 2020.10.26.525

Dengan Hasil sebagai Berikut:

Tingkat Kesamaan diseluruh artikel (*Similarity Index*) yaitu 4%

Demikian surat keterangan ini dibuat untuk digunakan sebagaimana mestinya.

Surabaya, 26 Oktober 2020

Ketua LPPM



UNUSA
LPPM

Dr. Ubaidillah Zuhdi, S.T., M.Eng., M.S.M.

NPP: 18101208

LPPM Universitas Nahdlatul Ulama Surabaya

Website : lppm.unusa.ac.id

Email : lppm@unusa.ac.id

Hotline : 0838.5706.3867

Monthly Water Balance Analysis for the Determination of Available

by

Submission date: 20-Oct-2020 12:00PM (UTC+0700)

Submission ID: 1420698108

File name: 15._Ketik_Ulang.PDF (824.26K)

Word count: 3450

Character count: 18216

BOOK 2


ISBN : 978-602-8915-93-9



ISNAR-C2FS 2011

International Seminar on Natural Resources, Climate Change and Food Security in Developing Countries

Proceeding





International Seminar Natural Resources, Climate Change and Food Security
in Developing Countries - ISNAR C2FS
June 27 - 28, 2011 Graha Pena Building, Surabaya - Indonesia
ISBN : 978-602-8915-93-9

Editorial Board

Alferd Hugo Volkaert, PhD. (CAB - University of Kasetsart, Thailand)

Zenaida Baoanan, PhD. (University of The Philippines Baguio, Philippines)

Katayama Takeshi, Prof. PhD. (University of Kagawa, Japan)

Sukendah, Dr. Ir. MSc. (University of Pembangunan nasional "Veteran" East Java, Indonesia)

Wani Hadi Utomo, Prof. PhD. (University of Tribuana, Malang, Indonesia)

Ricardo Bagarinao, PhD. (University of The Philippines Open University, Philippines)

Lin Qing, Prof. PhD. (School of Economic Fujian Normal University, China)

Surya Rathore, Prof. PhD. (University of Agriculture and Technology, Pantnagar, India)

Editor :

Yonny Koentjoro

Assistant Editor :

Wahyu Santoso

Pangesti Nugrahani

Makhziah

Cover :

Magnificent Digital Printing by : Purnomo Edi Sasongko

Proceeding design :

Yonny Koentjoro and Wahyu Santoso

Copyright : December 2011

Published by : Faculty of Agriculture, University of Pembangunan Nasional "Veteran" East Java, Indonesia

Held in between :

- Faculty of Agriculture, University of Pembangunan Nasional "Veteran" East Java, Indonesia
- Philippine Society for The Study of Nature (PSSN), Philippine
- CAB - University of Kasetsart, Thailand

Sub Theme : Agroforestry and Environmental Management
Oral Presentation

- | | | |
|-----|---|-----------|
| 31. | Carbon Pools and Spacial Variation of Carbon Sequestering Capability of Taba Penanjung Protection Forest, Bengkulu
Agus Susatya | 309 – 317 |
| 32. | Environmental Benefits and Costs of Conserving Layawan Watershed for Sustainable Domestic Water Supply in Oroquieta City, Philippines
Aisa O. Manlosa, Nicomedes D. Briones, Antonio J. Alcantara, Leonardo M. Florece, and Harold Glenn Valera | 318 – 330 |
| 33. | Monthly Water Balance Analysis for The Determination of Available Growing Season at North Samarinda District, East Kalimantan
Akas Pinaringan Sujalu and Akas Yekti Pulih Asih | 331 – 339 |
| 34. | Polluters and Water Quality of Cebu City River
Bonifacio S. Villanueva, Cecilio S. Baga and Corazon P. Macachor | 340 – 351 |
| 35. | Influence of Enzyme Treatment on The Quality of Cebu Technological University Deepwell Water
Cecilio S. Baga | 352 – 357 |
| 36. | Parasitoids And Their Effect On Teak Defoliator (Hyblaea Puera Crammer) In Teak Forest KPH Ngawi, East Java, Indonesia
Enggar Apriyanto, Edhi Martono, Sumardi, and Musyafa | 358 – 364 |
| 37. | Micro Watershed Model Development as Effort to Sustaining Water and Environment in Situbondo District
Evita Soliha Hani and Djoko Sudibya | 365 – 377 |
| 38. | Rice Waste Utilization and its Effects on Soil Properties in Selected Farms in Pangasinan, Philippines
Irene A. De Vera | 378 – 388 |
| 39. | Potent of Indonesian's Agricultural as Green House Gases Sources
K. A. Wijaya | 389 – 397 |
| 40. | The Impact of Motor Vehicle Density on Noise Level of Surabaya Main Roads
Naniek Ratni JAR and Erni Anita | 398 – 406 |
| 41. | Long-term Impact of Conventional Soil Management to Earthworm Diversity and Density on Sugarcane Plantation in East Java
Nurhidayati, E.Arisoesilarningsih, D.Suprayogo, and K.Hairiah | 407 – 419 |
| 42. | Drought Early Warning System And Preparation of Planting Pattern of Global Climate Changes in Connection With Mobile Technology-Based
Purnomo Edi Sasongko, I Gede Susrama MD, and M. Syahrul Munir | 420 – 427 |
| 43. | Prospects and Challenges in Harnessing Renewable Energy Source in the Philippines: the Case of Pico-hydro
Reynan P. Calderon, Gregorio J. Rodis, Adriano B. Singian, Julieta Laridad B. Reyes and Jose Paulo B. Tuazon | 428 – 435 |
| 44. | Proportion of Phenology of Arecaceae at Purwodadi Botanical Garden
Rony Irawanto | 436 – 445 |
| 45. | Nitrogen Cycling in Coffee Agroecosystems: N-Mineral and N-Leaching in the Presence of Different Litter Quality of Shade Trees
Priyadarshini, K.Hairiah, D. Suprayogo and J.B.Baon | 446 – 455 |
| 46. | Activity Change of Sulawesi Crested Black Macaques (Macaca Nigra) May Serve As An Early Indicator of Forest Degradation
Saroyo and Trina Tallei | 456 – 462 |
| 47. | Upland Rice Variety Assessment in Forest Areas Maintained Community together South Part in East Java
Sudarmadi Purnomo, Evy Latifah, Handoko and Sugiono | 463 – 473 |



**MONTHLY WATER BALANCE ANALYSIS FOR THE DETERMINATION
OF AVAILABLE GROWING SEASON AT NORTH SAMARINDA DISTRICT,
EAST KALIMANTAN**

Akas Pinarangan Sujalu¹ and Akas Yekti Pulih Asih²

¹ Agrotechnology Program Agriculture Faculty University of 17 Agustus 1945 Samarinda,
Jl. Ir. H. Juanda 80-Samarinda 75123

² Faculty of Economy, Kartini University -Surabaya
Jl Raya Nginden 19, Surabaya, Phone: 031-594-1954
Mobile Phone.085733030023

Mobile Phone. 081545695666: Email: pinaringan_b@yahoo.co.id

ABSTRACT

North Samarinda District this geographical area position 0°17'30"-0°30'00" LS dan 117°06'00"-117°22'00" BI, with the 31 475 Ha. Characterisfics of rain this area included in he Area Class III (1500 - 2000 mm/ year). With the Bimodel or Double Wave rainfall models with C patem. The hight rainfall depth periods at December and April, therefore the low rainfall depth at September and November. Have level Q = z 9.9 %, or rainfall tipe A (very wet area with tropical wet vegetation) and E1 agroclimte zone. Water Balance monthly indicated that this area have potential growing season about 12 months or all year round, have to water surplus 8 month as much as 478, 8 mm year' and water deficits about 3 months as much as 44.5 mm/year.

Keywords : Water Balance, Glowing Season

INTRODUCTION

Water is a natural resource that can be renewable and can be found everywhere, although 9till limited in quantity and quality of presence and availability both geographically and reviewed according to the season. Therefore, increased use will result in human intervention against the greater water resources. It will allow the change order and the hydrological cycle as more and more ten1tory and the presence of uneven distribution of water, both in spatial and temporal as well as decreasing watef quality. At the same time the utilization efficiency and lower water use and often ignore areas where water flow is derived, or river basin area (RBA).

Along with the development of the city, most of the upstream region of sub-watersheds have experienced pressure river Karangmumus degradation mainly due to logging, housing needs, and changes in regional function. Conditions are very obvious on a swamp area that stretches along the banks of the river downstream and the downstream Karangmumus sub-river of other rivers, a large part has been transformed into economic zones and settlements.



Management of sub-watershed with good manners and right is an attempt at controlling the reciprocal relationship between natural resources such as vegetation, soil and water by humans and all its activities. So that the basin management objectives to ensure sustainability and harmony of the ecosystem and increase the benefits of natural resources contained therein for human life can be achieved. It is intended as an effort to maintain the Mahakam river water discharge in the dry season which can prevent the intrusion of sea water. It also avoids increasing the flow rate of surface water in the rainy season which can lead to the high frequency of flooding in the area of the city of Samarinda (Trisusanto, 2002).

The direct impact that is felt is the change in the basin water balance Karangmumus tangible with the occurrence of droughts and floods are more widespread and more frequently lashed various aspects of life (Suyitno, 1989 . Anonymous, 2001).

MATERIALS AND METHOD

a. time and Place Research

The study was conducted at North Samarinda District for approximately 6 (six) months (July-December 2008) in an area of approximately ± 31 475 hectares.

b. Data Collection

Data collected from both primary and secondary data related to the research include:

- a. Climate, particularly rainfall and evaporation
- b. Physiographic characteristics, particularly land slope
- c. Soil conditions, especially those related to water status in soil
- d. The vegetation, especially the dominance of vegetation, land cover

c. Water Balance Analysis

Analysis of water balance of land stated in the form of integral equations by simplifying some similarities, so that the water balance of a land area can be expressed in the form of the equation:

$$CH = \text{ETA} \pm \Delta \text{WCS} \pm \text{Li}$$

Where: CH = rainfall (mm months⁻¹)

ETA = actual evapotranspiration (SETP)

ΔWCS = soil water content changes (mm months⁻¹)

Li = runoff (surplus or deficit depending on its value) (mm month)



c.1*. analysis of Potential Evapotranspiration (ETP)

Calculation of potential evapotranspiration (ETP) using equations from Buckman and Braddy (1969), quoted by Sujalu (1997), as follows:

$$EPT_i = 616X \left(10X \frac{T_i^a}{T} \right)$$

$$I = \sum_{jan}^{des} \left(\frac{T_i}{5} \right)^{1.514}$$

$$8 = 6.75 \times 10^{-1} I^* - 7.71 \times 10^{-2} I^{*2} + 1.792 \times 10^{-1} I + 0.492$$

where; ETP = Evapotranspiration

T_i = temperature of the month to the first monthly

I = Index monthly heat

a = Constant

C.2. analysis of Soil Water Content (WCS)

Changes in Water Content of Soil (WCS) is the difference in soil moisture content on a period to prior periods between sequential. For each change in soil water content, can be calculated with the formula R - ETP that if a negative value, there will be a deficit (lack of) water for (ETP = Eta). Conversely, if (R - ETP) is positive, then there will be a surplus/excess of water (R-ETP-AWCS), so that soil water availability decreases water exponentially and expressed by the equation:

$$SWA = WHC \times k^a,$$

$$WHC - FC - PWP,$$

$$WCS = PWP + ASW$$

Where:

WCS = Actual Soil Water Content (mm)

ASW = Availability of Soil Water Actual (mm)

WHC = water holding capacity or availability of Maximum Soil Water (mm)

FC = Field Capacity (mm)

PWP = Permanent Wilt Point (mm)

K = Constant (obtained $k = ((P_0 + P_1) / WHC)$),

with

P₀ = 1.000412351 and P₁ = -1.073807306

a = accumulation of potentially lost water (Accumulate Potential Water)



Loss, APWL), which represents the accumulated value (R-ETP) when the value of R < ETP

RESULTS AND DISCUSSION

A. Preview Area North-Samarinda District

North-Samarinda District geographically located in the region North Samarinda district is part of the Mahakam river basin is located at coordinates between 0 17'30"-0 30'00" SL and 117 06'00"-117 22'00" EW with a total area reaches 31 475 hectares. This area is traversed by two creeks ie. sub-watershed areas, namely sub-sub watershed Karangmumus and sub-sub-watershed Siring as well as several other small rivers.

North Samarinda district has the largest area, 20,520 hectares, or 28.58 A of total large city. Similarly, the total population, 138,726 people. Generally inhabiting North Samarinda is transmigran district of Java, is a district with the most extensive region in Samarinda and herein lies Siring River Airport is under construction. North Samarinda district includes 6 (five) sub-districts, namely sub-district Le ake, Pelita, sempaja utara, Sempaja Selatan, Sungai Siring and Tanah Merah.

This area has varied topography, with elevation ranging from topographic region 10-120 m above sea level with a diverse variety of heights.

Table 1. The total area of the basin based on gradient class North Samarinda district

The range of slope (%)	Slope Class Size	Area	
		Hectares	(%)
0 ~ 8	I (Flat)	4.907,85	15,59
8 ~ 15	II ((flat)(Landai)	2.780,66	8,83
15 ~ 25	III (Rather Steep)	18.134,51	57,62
25 ~ 40	IV (Steep)	6.398,13	20,33
>40	V (Very Steep)	472,93	1,50
□ amount		31.475	100.00

Source : Anonim (2001)



Appendix Table 1. Monthly Water equilibrium in the North Samarinda District area
 (0°17'30"- 0°30'00" SL and 117°06'00"-117°22'00" EL)

Parameter	Month											
	Jan	Feb	Mar	Apr	May	June	July	Agst	Sept	Okc	Nov	Des
Rainfall (mm)	194.0	143.0	233.0	333.0	183.0	113.0	178.0	121.0	104.0	134.0	198.0	214.0
Potential Evapotranspiration (mm)	139.0	138.7	139.6	140.2	139.6	138.4	137.2	138.1	138.4	139.3	139.3	138.7
R - ETP (mm)	55.0	4.3	94.4	192.8	42.4	-25.4	40.8	-17.1	-34.4	-5.3	58.7	75.3
APWL (mm)	0	0	0	0	0	0	0	-17.1	-51.5	-61.8	0	0
Water Content of Soil WCS (mm) ¹	268	268	268	268	268	242.6	268	250.9	198.5	136.7	185.4	260.7
ΔWCS	0	0	0	0	0	-25.8	0	-17.1	-58.0	-62.0	-3.3	0
Actual Evapotranspiration/ETA (mm) ²	139.0	138.7	139.6	140.2	139.6	138.4	136.9	142.0	152.0	196.0	139.3	138.7
Deficit (mm)	0	0	0	0	0	0	0.3	3.9	13.6	26.7	0	0
Surplus (mm)	55.0	4.3	94.4	192.8	42.4	0	0	0	0	0	58.7	75.3

Description: 1. water content of soil (WCS) at Field Capacity (FC)

2. Actual evapotranspiration (ETA) in the period of time deficit (R<ETP) was obtained from R (mm) + □ WCS. While at the time of surplus (R> ETP) the amount equal to ETP

Table 2. The Area Land Use Type at North Samarinda district

No.	The Area Land Use Type	Area	
		(ha)	(%)
01.	Farm (dry land farming)	203,13	0,65
02.	Forest	146,15	0,46
03.	Shrub	6.996,25	22,23
04.	Mixed Garden	4.473,44	14,21
05.	Bush	14.501,36	46,07
06.	Wetland	648,99	2,06
07.	Garden	1.106,64	3,52
08.	Settlements	2.267,78	7,21
09.	Settlement expansion (Pp)	215,61	0,69
10.	Slough/swamp area	915,63	2,91
Amount		31.475,00	100

Source : Anonim (2001)

Table 3. Climate Data Average Monthly North Samarinda district area
 (0°17'30"- 0°30'00" SL and 117°06'00"-117°22'00" EL)

Climate Elements	Month											
	1	2	3	4	5	6	7	8	9	10	11	12
Rainfall (mm month ⁻¹)	194	123	233	333	183	113	178	121	104	134	198	214
Rainy Days (days)	12	11	12	14	12	11	9	10	9	11	11	12
Temperature (°C)	26.8	26.7	27.0	27.2	27.0	26.6	26.2	26.5	26.6	26.9	26.9	26.9
Humidity (%)	87.2	86.3	89.2	90.2	88.6	86.5	85.6	86.2	83.4	85.9	87.1	86.3
Sun Radiation (Kkal cm ⁻²)	0.55	0.48	0.51	0.53	0.53	0.51	0.51	0.49	0.41	0.44	0.44	0.52



B. Condition Elements The climate in the North Samarinda district

Based on rainfall data from 3 (three) climate observation station in the basin area Karangmumus year period from 2001 to 2009 showed that rainfall monthly average ranged from 104-214 mm month⁻¹ or an average of 168 mm month⁻¹, whereas The average rainfall ranging from 1510-2850 mm year¹ or average of 2018 mm year¹. Rainfall occurred on rainy days (rd) monthly rates ranging from 9-14 rd with an average rainfall occurred 11 rd month["] .

Analysis of rainfall characterization includes four main components, namely:

1. Annual Rainfall spread of this area falls within Class Region III (rainfall between 1500-2000 mm year¹).
2. Spread Type Rainfall North Samarinda district area has a period of dry months (months with rainfall of <100 mm month^{"1}). Thus obtained value of $Q = -J-9.996$, or rain type A (which may imply that North Samarinda district is very wet areas with dense vegetation of tropical rain
3. Rainfall patterns or Bimodel Dual (Double Wave) with the notation Pattern C, periods of high rainfall occurred in December and April, while periods of low rainfall occurred in September and November.
4. Agro-climate zones, the North Samarinda district the same as the basin area Karangmumus has a 1 (one) dry months (DM), 8 months humid (HM) and three wet months (WM), including agro-climate zones E1.

C. Calculation of Potensial Evapotranspiration (ETP)

Calculation of potential evapotranspiration (ETP) conducted monthly by using equations from Buckman and Braddy (1969), quoted by Sujalu (1997) in the North Samarinda district ranged from 137.2-140.2 mm or an average of 138.8 mm month⁻¹ the highest point in months April amounted to 140.2 mm and the lowest points in July amounted to 137.2 mm.

D. Water Balance

The calculation result in soil water status wewers obtained from analysis of soil physical properties in the laboratory soil Assessment Institute for Agricultural Technology (BPTP) East Kalimantan Prov. from Heriansyah (2004) showed that soil available water content (WCS) in the range between 244-299mm or average 268 mm

Water equilibrium implies about the details of the input and outputs of water in one place at a certain time period, compiled in the form of quantitative equations,



which provide information in the form of quantitative values of each component of input and output water, can be seen in Appendix Table 1.

Monthly Water Equilibrium analysis of the results mentioned above can be seen that these areas have a surplus during the eight months that occurred in a period of months from January to June and in November-December. The monthly water surplus in detail is in January amounted to 27.0 mm, 57.3 mm in February amounted, in March amounted 119.7 mm (the highest monthly surplus), in April amounted to 72.6 mm, 48.4 mm in May, months of June amounted to 19.6 mm (the lowest monthly surplus), the month of November amounted to 58.7 months in December and amounted to 75.3 mm in overall water surplus reached 478.8 mm year. In addition to having monthly water surpluses, the region normally monthly cumulative water deficit in a period of months from June to October as a whole as much as 44.5 mm year⁻¹, with details of the deficit in June amounted to 0.4 mm month⁻¹, July amount 0.3 mm month⁻¹, the month of August amounted to 3.9 mm month⁻¹, the month of September amounted to 13.6 mm month⁻¹ and in October of 26.7 mm month⁻¹.

As has been previously communicated its position Karangmumus river divides the city of Samarinda especially North Samarinda District, and considering the amount of potential run-off that occurred in the region and also by considering the conditions Karangmumus area topography, the basin area Karangmumus at the North Samarinda Regions very possible to build dams or reservoirs, which have various functions. Although the main function is to accommodate the construction of the dam monthly surplus water run off resulting in the potential is big enough in this area, as well as water reserves in the period in the months of water deficit that can be utilized by a variety of purposes including drinking water.

E. Analysis of Cropping Periods (Growing Season)

To determine the length of cropping period (the length of growing season) can be done based on the ratio P/PE (ratio between precipitation and potential evapotranspiration), defined as the time interval in a year that have a ratio P/PE > 0.5 plus the time needed for evapotranspiration 100 mm of ground water is considered available in the soil (FAO, 1978). Results of analysis ratio P/PE can be seen in Table 4 below.



Table 4. Ratio rainfall (R) and potential evapotranspiration (EP) monthly

Climate Elements	Months											
	1	2	3	4	5	6	7	8	9	10	11	12
Rainfall (mm/month)	194	123	233	333	183	113	178	121	104	134	198	214
Evapotr.Pot. / ETP (mm)	139.0	138.7	139.6	140.2	139.6	138.4	137.2	138.1	138.4	139.3	139.3	138.7
Ratio P/PE	1.4	0.9	1.7	2.4	1.3	0.8	1.3	0.9	0.8	0.9	1.4	1.5

Based on this analysis the ratio P / PE ratio of the above in mind that the P / PE in the rain fall average monthly cumulative throughout the year (12 months) is always» 0.5. Therefore, according to the restrictions provided FAO (1978), the North Samarinda District afeas have planting period (the length of growing season) for 12 months or all year round.

CONCLUSION

Based on the description as a whole can be concluded that North Samarinda districtarea has a all year round (12 month) potential planting period (growing season) which is supported by the surplus water during 7 (seven) months or cumulatively amounted to 478.8 mm year-1 and a deficit of water for four months or cumulative 44, 5 mm years-1.

REFERENCES

- Anonymous. 2001. Inventory and Identification of Areas Prone Environment in Sub-watershed Karangmumus Samarinda (Mapping erosion hazard level). BAPEDALDA. East Kalimantan.
- Asdak, C. 1995. Hydrology and Watershed Management. Gadjah Mada University Press. Yogyakarta.
- Sujalu, A.P. 1997. Climate Resources Utilization for Agriculture and Water Management Planning. Seminar Papers. LPTP-East Kalimantan.
- . 2000. Water Balance Analysis of Land in Pasir District. Journal of Agriculture Vol. 6 No. 2 Septembr 2000 - Faculty of Agriculture Mulawarman University
- — 2002. Rice Cropping Pattern Planning, com and soybeans on 60•â Chance Rain Events in Tenggaraong - Xutai Regency. HABITAT Journal Vol. No. XIII. June 2, 2002, Faculty of Agriculture. Brawijaya University
- Seyhan, E. 1990. Hydrology Basics. Gadjah Mada University Press. Yogyakarta.



Sosrodarsono, S. and Takeda. 1983. Hydrology for Water Resources. PT Pradnya Paramitha. Jakarta.

Trisusanto, A. 2002. Protected Area Management Plan for the Upper Rock watershed Mumus Samarinda through Local Community approach. Thesis. Graduate Program. Faculty of Civil Engineering. Surabaya Institute of Technology. (Unpublished)



15._Ketik_Ulang.PDF

ORIGINALITY REPORT

4%

SIMILARITY INDEX

3%

INTERNET SOURCES

0%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

1

Submitted to UPN Veteran Jawa Timur

Student Paper

2%

2

repository.unusa.ac.id

Internet Source

1%

3

jesam-uplb.org

Internet Source

1%

4

repo.unsrat.ac.id

Internet Source

1%

Exclude quotes On

Exclude bibliography On

Exclude matches < 1%