



# **ICSAFS**

**International Conference  
on Sustainable Agriculture and Food Security:  
*Challenges and Opportunities***

## **Proceeding**

**Universitas Padjadjaran  
Indonesia  
2012**



# **ICSAFS**

**International Conference  
on Sustainable Agriculture and Food Security:  
*Challenges and Opportunities***

## **Proceeding**

**(Poster Papers)**

Editors:

Anne Nurbaity (Indonesia)  
Edy Subroto (Indonesia)  
Endang Yuni Setyowati (Indonesia)  
Florin Stanica (Romania)  
Ichsan Nurul Bari (Indonesia)  
Klaus Wimmers (Germany)  
Nono Carsono (Indonesia)  
Oviyanti Mulyani (Indonesia)  
Pasi Lehmousloto (Finlandia)  
Paul S. Teng (Singapore)  
Shantosa Yudha Siswanto (Indonesia)  
Stevica Aleksic (Republic of Serbia)

**Universitas Padjadjaran**

**Indonesia  
2012**

## DISCLAIMER

The International Conference on Sustainable Agriculture and Food Security (ICSAFS) was held on September 2011 at Universitas Padjadjaran. Responsibility for the content of these papers rests solely with the authors. Where trade names are used, this constitutes neither endorsement of nor discrimination against any product by the institution.

Published by Faculty of Agriculture Universitas Padjadjaran Indonesia

April 2012

Nurbaity, A., Subroto, E., Setyowati, E.Y., Stanica, F., Bari, I.N., Carsono, N., Mulyani, O., Lehmousloto, P., Teng, P.S., Siswanto, S.Y., Aleksic, S. 2012. Proceeding of International Conference on Sustainable Agriculture and Food Security (ICSAFS).

Copyright on all papers on the Conference resides with Universitas Padjadjaran  
Jl. Raya Jatinangor km. 21 Bandung 40600 West Java Indonesia

ISBN 978-9798426-13-5 [printed version]

ISBN 978-979-8246-14-2 [electronic version]

Technical editing and design: Ichsan Nurul Bari and Gigih Ibnu Prayoga

# Water Balanced Analysis to Available Growing Season at Sungai Pinang District-East Kalimantan

Sujalu, A. P.

Faculty of Agriculture, 17<sup>th</sup> August 1945 University of Samarinda,

Jl. Ir. H. Juanda 80 - Samarinda 75123

E-mail: pinaringan\_b@yahoo.co.id

## Abstract

The geographical position of Sungai Pinang District are between  $0^{\circ}17'30''$ - $0^{\circ}30'00''$  South Latitude and  $117^{\circ}06'00''$ - $117^{\circ}22'00''$  East Longitude, with the land area is 34,165 Ha. Characteristics of rain in this area are in the Area Class III (1500 – 2000 mm/year). With the Bimodel or Double Wave rainfall models with C pattern. The high rainfall period is on December to April, and the low rainfall is on September to November. The level  $Q = \pm 9.2\%$ , or rainfall type A (very wet area with tropical wet vegetation) and E1 agroclimate zone. Water Balance monthly indicated that this area have potential growing season about 9 months, have to water surplus 8 month ( $472.6 \text{ mm year}^{-1}$ ) and water deficits about 3 months ( $44.5 \text{ mm/year}$ ).

**Keywords:** water balanced, growing season

## Introduction

Water is a natural resource that can be renewable and can be found everywhere, although still limited in quantity and quality of presence and availability both geographically and reviewed according to the season. Therefore, increased use will result in human intervention against the greater water resources. It will allow the change order and the hydrological cycle as more and more territory and the presence of uneven distribution of water, both in spatial and temporal as well as decreasing water quality. At the same time the utilization efficiency and lower water use and often ignore areas where water flow is derived, or river basin area (RBA).

Along with the development of the city, most of the upstream region of sub-watersheds has experienced pressure river Karangmumus degradation mainly due to logging, housing needs, and changes in regional function. Conditions are very obvious on a swamp area that stretches along the banks of the river downstream and the downstream Karangmumus sub-river of other rivers, a large part has been transformed into economic zones and settlements.

Management of sub-watershed with good manners and right is an attempt at controlling the reciprocal relationship between natural resources such as vegetation, soil and water by humans and all its activities. So that the basin management objectives to ensure sustainability and harmony of the ecosystem and increase the benefits of natural resources contained therein for human life can be achieved. It is intended as an effort to maintain the Mahakam river water discharge in the dry season which can prevent the intrusion of sea water. It also avoids increasing the flow rate of surface water in the rainy season which can lead to the high frequency of flooding in the area of the city of Samarinda (Trisusanto, 2002).

The direct impact that is felt is the change in the basin water balance Sungai Pinang District with the occurrence of droughts and floods are more widespread and more frequently lashed various aspects of life (Suyitno, 1989; Anonymous, 2001).

## Methods

### a. Time and place research

The study was conducted at Sungai Pinang District for approximately 6 (six) months (July-December 2008) in an area of approximately  $\pm 31,475$  hectares.

### b. Data collection

Data collected from both primary and secondary data related to the research, include:

- Climate, particularly rainfall and evaporation
- Physiographic characteristics, particularly land slope
- Soil conditions, especially those related to water status in soil
- The vegetation, especially the dominance of vegetation, land cover

### c. Water balance analysis

Analysis of water balance of land stated in the form of integral equations by simplifying some similarities, so that the water balance of a land area can be expressed in the form of the equation:

$$CH = ETA \pm \Delta WCS \pm Li$$

Where:

CH = rainfall (mm months<sup>-1</sup>)

ETA = actual evapotranspiration ( $\leq$ ETP)

$\Delta WCS$  = soil water content changes (mm months<sup>-1</sup>)

Li = runoff (surplus or deficit depending on its value) (mm months<sup>-1</sup>)

#### C.1. Analysis of Potential Evapotranspiration (ETP)

Calculation of potential evapotranspiration (ETP) using equations from Buckman and Braddy (1969), quoted by Sujalu (1997), as follows:

$$EPT_i = 616 \times \left( 10 \times \frac{T_i}{I} \right)^{a1}$$

$$I. = \sum_{jan}^{des} \left( \frac{T_i}{5} \right)^{1.514}$$

$$a = 6.75 \times 10^{-7} I^3 - 7.71 \times 10^{-5} I^2 + 1.792 \times 10^{-2} I + 0.492$$

Where;

ETP = Evapotranspiration

Ti = temperature of the month to the first monthly

I = Index monthly heat

a = Constant

## C.2. Analysis of Soil Water Content (WCS)

Changes in Water Content of Soil (WCS) is the difference in soil moisture content on a period to prior periods between sequential. For each change in soil water content, can be calculated with the formula  $R - ETP$  that if a negative value, there will be a deficit (lack of) water for ( $ETP = \text{Eta}$ ). Conversely, if  $(R - ETP)$  is positive, then there will be a surplus/excess of water ( $R - ETP - \Delta WCS$ ), so that soil water availability decreases water exponentially and expressed by the equation:

$$\begin{aligned} ASW &= WHC \times k^a, \\ WHC &= FC - PWP, \\ WCS &= PWP + ASW \end{aligned}$$

Where:

WCS = Actual Soil Water Content (mm)

ASW = Availability of Soil Water Actual (mm)

WHC = water holding capacity or availability of Maximum Soil Water (mm)

FC = Field Capacity (mm)

PWP = Permanent Wilt Point (mm)

k = Constant (obtained  $k = ((P_0 + P_1) / WHC)$ ,

With

$P_0 = 1.000412351$  and  $P_1 = -1.073807306$

a = accumulation of potentially lost water (Accumulate Potential Water Loss, APWL), which represents the accumulated value  $(R - ETP)$  when the value of  $R < ETP$

## Results and Discussion

### A. Preview area of sungai pinang district

Sungai Pinang District geographically located in the region of the Mahakam watershed is located at coordinates between  $0^{\circ}17'30'' - 0^{\circ}30'00''$  SL and  $117^{\circ}06'00'' - 117^{\circ}22'00''$  EW with a total area reaches 34165 hectares. This area is traversed by two sub-sub watershed Karangmumus and sub-sub-watershed Siring as well as several other small rivers.

Sungai Pinang District has the largest area, Samarinda city about 29.78% of total large city. Similarly, the total population, 138,726 people. Generally inhabiting this district is transmigran district of Java, is a district with the most extensive region in Samarinda and herein lays Siring River Airport or New Airport of Samarinda is under construction. Sungai Pinang District includes 6 (five) sub-districts, namely sub-district Lempake, Pelita, Sempaja Utara, Sempaja Selatan, Sungai Siring and Tanah Merah, and has varied topography, with elevation ranging from topographic region 10-120 m.asl with a diverse variety of heights.

Table 1. The total area of the basin based on gradient class Sungai Pinang District

The range of slope (%)	Slope Class Size	Area	
		Hectares	(%)
0 ~ 8	I (Flat)	4907.85	15.59
8 ~ 15	II ((flat)(Landai)	2780.66	8.83
15 ~ 25	III (Rather Steep)	18134.51	57.62
25 ~ 40	IV (Steep)	6398.13	20.33
>40	V (Very Steep)	472.93	1.50
Amount		34165.00	100.00

Sumber : Anonim (2009)

Table 2. The Area Land Use Type at Sungai Pinang District

No.	The Area Land Use Type	Area	
		(ha)	(%)
01.	Farm (dry land farming)	503.13	0.65
02.	Forest	146.15	0.46
03.	Shrub	6996.25	22.23
04.	Mixed Garden	5473.44	14.21
05.	Bush	15501.36	46.07
06.	Wetland	648.99	2.06
07.	Garden	1106.64	3.52
08.	Settlements	2267.78	7.21
09.	Settlement expansion (Pp)	215.61	0.69
10.	Slough/swamp area	915.63	2.91
Amount		34165.00	100.00

Sumber : Anonim (2009)

### B. The Elements of Climate Condition in the Sungai Pinang District

Based on rainfall data from 3 (three) climate observation station in the Karangmumus watershed year period from 2001 to 2009 showed that rainfall monthly average ranged from 103-216 mm month<sup>-1</sup> or an average of 171 mm month<sup>-1</sup>, whereas the average rainfall ranging from 1515-2565 mm year<sup>-1</sup> or average of 2028 mm year<sup>-1</sup>. Rainfall occurred on rainy days (rd) monthly rates ranging from 9-14 rd with an average rainfall occurred 11<sup>rd</sup> month<sup>-1</sup>.

Analysis of rainfall characterization includes four main components, namely:

1. Annual Rainfall spread of this area falls within Class Region III (rainfall between 1500-2000 mm year<sup>-1</sup>).
2. Spread Type Rainfall Sungai Pinang District area has a period of dry months (months with rainfall of <100 mm month<sup>-1</sup>). Thus obtained value of Q = ±9.2%, or rainfall type A (which may imply that Sungai Pinang District is very wet areas with dense vegetation of tropical rain).
3. Rainfall patterns or Bimodel Dual (Double Wave) with the notation Pattern C, periods of high rainfall occurred in December and April, while periods of low rainfall occurred in September and November.
4. Agro-climate zones, the Sungai Pinang District the same as the Karangmumus watershed has a 1 (one) dry months (DM), 8 months humid (HM) and three wet months (WM), including agro-climate zones E1.

### C. Calculation of potential evapotranspiration (ETP)

Calculation of potential evapotranspiration (ETP) conducted monthly by using equations from Buckman and Braddy (1969), quoted by Sujalu (1997) in the Sungai Pinang District ranged from 137.2-140.2 mm or an average of 138.8 mm month<sup>-1</sup>, the highest point in months April amounted to 140.2 mm and the lowest points in July amounted to 137.2 mm.

Table 3. Monthly Climate Data Average Sungai Pinang District Area (0°17'30"- 0°30'00" SL and 117°06'00"-117°22'00" EL)

Climate Elements	Month											
	1	2	3	4	5	6	7	8	9	10	11	12
Rainfall (mm month <sup>-1</sup> )	194	123	233	333	183	113	178	121	104	134	198	214
Rainy Days (days)	12	11	12	14	12	11	9	10	9	11	11	12
Temperature (°C)	26.8	26.7	27.0	27.2	27.0	26.6	26.2	26.5	26.6	26.9	26.9	26.9
Humidity (%)	87.2	86.3	89.2	90.2	88.6	86.5	85.6	86.2	83.4	85.9	87.1	86.3
Sun Radiation (Kkal cm <sup>-2</sup> )	0.55	0.48	0.51	0.53	0.53	0.51	0.51	0.49	0.41	0.44	0.44	0.52

#### D. Water balance

The calculation result in soil water status were obtained from analysis of soil physical properties in the Laboratory Soil Assessment Institute for Agricultural Technology (BPTP) East Kalimantan Province from Heriansyah (2004) showed that soil available water content (WCS) in the range between 247-297mm or average 268 mm

Water equilibrium implies about the details of the input and outputs of water in one place at a certain time period, compiled in the form of quantitative equations, which provide information in the form of quantitative values of each component of input and output water, can be seen in Appendix Table 1.

Monthly Water Equilibrium analysis of the results mentioned above can be seen that these areas have a surplus during the eight months that occurred in a period of months from January to June and in November-December. The monthly water surplus in detail is in January amounted to 27.0 mm, February amounted 57.3 mm, in March amounted 119.7 mm (the highest monthly surplus), in April amounted to 72.8 mm, 48.4 mm in May, months of June amounted to 19.6 mm (the lowest monthly surplus), the month of November amounted to 58.7 months in December and amounted to 75.3 mm in overall water surplus reached 476.2 mm year<sup>-1</sup>.

In addition to having monthly water surpluses, the region normally monthly cumulative water deficit in a period of months from June to October as a whole as much as 44.5 mm year<sup>-1</sup>, with details of the deficit in June amounted to 0.4 mm month<sup>-1</sup>, July amount 0.3 mm month<sup>-1</sup>, the month of August amounted to 3.9 mm month<sup>-1</sup>, the month of September amounted to 13.6 mm month<sup>-1</sup> and in October of 26.7 mm month<sup>-1</sup>.

As has been previously communicated its position Karangmumus river divides the city of Samarinda especially Sungai Pinang District, and considering the amount of potential run-off that occurred in the region and also by considering the conditions Karangmumus area topography, the basin area Karangmumus at the North Samarinda Regions very possible to build dams or reservoirs, which have various functions. Although the main function is to accommodate the construction of the dam monthly surplus water run off resulting in the potential is big enough in this area, as well as water reserves in the period in the months of water deficit that can be utilized by a variety of purposes including drinking water



### E. Analysis of cropping periods (growing season)

To determine the length of cropping period (the length of growing season) can be done based on the ratio P/PE (ratio between precipitation and potential evapotranspiration), defined as the time interval in a year that have a ratio P/PE>0.5 plus the time needed for evapotranspiration 100 mm of ground water is considered available in the soil (FAO, 1978). Results of analysis ratio P/PE can be seen in Table 4 below.

Table 4. The Ratio of Rainfall (R) and Potential Evapotranspiration (EP) Monthly

Climate Elements	Months											
	1	2	3	4	5	6	7	8	9	10	11	12
Rainfall (mm / month)	194	123	233	333	183	113	178	121	104	134	198	214
Evapotr.Pot. / ETP (mm)	139.0	138.7	139.6	140.2	139.6	138.4	137.2	138.1	138.4	139.3	139.3	138.7
Ratio P/PE	1.4	0.9	1.7	2.4	1.3	0.8	1.3	0.9	0.8	0.9	1.4	1.5

Based on this analysis the ratio P / PE ratio of the above in mind that the P / PE in the rain fall average monthly cumulative throughout the year (12 months) is always> 0.5. Therefore, according to the restrictions provided FAO (1978), the Sungai Pinang District areas have planting period (the length of growing season) for 12 months or all year round.

### Conclusion

Based on the description as a whole can be concluded that Sungai Pinang District area has a all year round (12 month) potential planting period (growing season) which is supported by the surplus water during 8 (seven) months or cumulatively amounted to 476.2 mm year-1 and a deficit of water for 3 months or cumulative 44, 5 mm years<sup>-1</sup>.

### Reference

- Anonymous. 2009. Inventory and Identification of Areas Prone Environment in Sub-watershed Karangmumus Samarinda (Mapping erosion hazard level). BAPEDALDA. East Kalimantan.
- Asdak, C. 1995. Hydrology and Watershed Management. Gadjah Mada University Press. Yogyakarta.
- Sujalu, A.P. 1997. Climate Resources Utilization for Agriculture and Water Management Planning. Seminar Papers. LPTP-East Kalimantan.
- . 2000. Water Balance Analysis of Land in Pasir District. Journal of Agriculture Vol. 6 No. 2 September 2000 - Faculty of Agriculture. Mulawarman University
- . 2002. Rice Cropping Pattern Planning, corn and soybeans on 60% Chance Rain Events in Tenggaraong – Kutai Regency. HABITAT Journal Vol. No. XIII. June 2, 2002, Faculty of Agriculture. Brawijaya University
- Seyhan, E. 1990. Hydrology Basics. Gadjah Mada University Press. Yogyakarta.
- Sosrodarsono, S. and Takeda. 1983. Hydrology for Water Resources. PT Pradnya Paramitha. Jakarta.
- Trisusanto, A. 2002. Protected Area Management Plan for the Upper Rock watershed Mumus Samarinda through Local Community approach. Thesis. Graduate Program. Faculty of Civil Engineering. Surabaya Institute of Technology. (Unpublished)